Zero temperature black holes in semiclassical gravity

Paul R. Anderson*
Department of Physics, Wake Forest University, Winston-Salem, NC 27109, USA

William A. Hiscock†
Department of Physics, Montana State University, Bozeman, MT 59717, USA

Brett E. Taylor‡
Department of Chemistry and Physics, Radford University, Radford, VA 24142, USA

arXiv:gr-qc/0102111v3 14 Mar 2001

Abstract

The semiclassical Einstein equations are solved to first order in $\epsilon = \hbar/M^2$ for the case of an extreme or nearly extreme Reissner-Nordström black hole perturbed by the vacuum stress-energy of quantized free fields. It is shown that, for realistic fields of spin 0, 1/2, or 1, any zero temperature black hole solution to the equations must have an event horizon at $r_h < |Q|$, with Q the charge of the black hole. It is further shown that no black hole solutions with $r_h < |Q|$ can be obtained by solving the semiclassical Einstein equations perturbatively.

Static spherically symmetric zero temperature black holes have proven to be very interesting and important at the classical, semiclassical, and quantum levels. Classically the only static spherically symmetric black hole solution to Einstein's equations with zero surface gravity (and hence zero temperature) is the extreme Reissner-Nordström (ERN) black hole, which possesses a charge equal in magnitude to its mass. At the quantum level, the statistical mechanical entropy of zero temperature (extreme) black holes has been calculated in string theory [1] and shown to be identical to the usual Bekenstein-Hawking formula for the thermodynamic entropy. The usual semiclassical temperature and entropy calculations for ERN black holes have all been made in the test field approximation where the effects of quantized fields on the spacetime geometry are not considered. However, it is well known that quantum effects alter the spacetime geometry near the event horizon of a black hole. In particular they can change its surface gravity and hence its temperature [2–4].

In a previous paper [5] we examined the semiclassical backreaction due to the vacuum stress-energy of massless and massive free quantized fields with spin 0, 1/2, and 1 on a static Reissner-Nordström (RN) black hole. We calculated the first-order (in $\hbar$) corrections to the RN background and found, in all cases examined, that the surface gravity was nonzero because the energy density of the quantized field on the horizon is negative, $\langle T^t_t \rangle > 0$. [6] This led us to conclude that macroscopic zero temperature black holes might not exist in semiclassical gravity. After this work was published a paper appeared by Lowe [7] which finds a zero temperature solution to the linearized semiclassical backreaction equations.

In this paper we investigate the perturbation series more generally than we did previously [5]. We find that for nearly extreme black holes, the surface gravity (and temperature) of the black hole decreases as the radius of the event horizon r_h decreases. Thus these solutions form a sequence and the sequence appears to terminate at the zero temperature solution found by Lowe [7]. In our previous paper the lowest temperature solution examined had $r_h = |Q|$ and $\kappa = 4\pi|Q|\langle T_t^t\rangle > 0$; Lowe's proposed solution has $r_h = |Q|(1-4\pi Q^2\langle T_t^t\rangle) < |Q|$ and $\kappa = 0$. We discuss the validity of the solutions in this sequence and show that it is not possible to use perturbation theory to obtain those with $r_h < |Q|$, including the one proposed by Lowe. However it is possible that they nevertheless approximate exact solutions to the full nonlinear semiclassical equations. We also discuss a potential problem that may occur for all zero temperature black hole solutions to the semiclassical backreaction equations.

The general static spherically symmetric metric can be written in the form [6]:

$$ds^2 = -f(r)dt^2 + h(r)dr^2 + r^2 d\Omega^2 \ , \tag{1}$$

where $d\Omega^2$ is the metric of the two-sphere. The metric can describe a black hole with an event horizon at $r = r_h$ if $f(r_h) = 0$. To avoid having a scalar curvature singularity at the event horizon it is necessary that $h^{-1}(r_h) = 0$ as well. The surface gravity of such a black hole is

$$\kappa = \left(\frac{1}{2}\right) \frac{f'}{\sqrt{fh}} \Bigg|_{r=r_h} \ , \tag{2}$$

where the prime represents a derivative with respect to r and the expression is evaluated at the horizon radius, r_h. The temperature is then $T = \kappa/(2\pi)$.

Since we wish to perturb the spacetime with the vacuum energy of quantized fields, we begin by considering the general Reissner-Nordström metric as the "bare" state. For the RN metric,

$$f(r) = h^{-1}(r) = 1 - \frac{2M}{r} + \frac{Q^2}{r^2} \ , \tag{3}$$

where Q is the electric charge and M is the mass of the black hole. The outer event horizon is located at

$$r_+ = M + \sqrt{M^2 - Q^2} \ . \tag{4}$$

For the ERN black hole $|Q| = M$.

In semiclassical gravity, the geometry is treated classically while the matter fields are quantized. In examining the semiclassical perturbations of the RN metric caused by the vacuum energy of quantized fields, we continue to treat the background electromagnetic field as a classical field. The right hand side of the semiclassical Einstein equations will then contain both classical and quantum stress-energy contributions,

$$G_{\mu\nu} = 8\pi \left[(T_{\mu\nu})^C + \langle T_{\mu\nu} \rangle \right] \ . \tag{5}$$

We consider the situation where the black hole is in thermal equilibrium (whether at zero or nonzero temperature) with the quantized field; the perturbed geometry then continues

to be static and spherically symmetric. To first order in $\epsilon = \hbar/M^2$ the general form of the perturbed RN metric may be written as:

$$ds^2 = -[1 + 2\epsilon\rho(r)]\left(1 - \frac{2m(r)}{r} + \frac{Q^2}{r^2}\right)dt^2 + \left(1 - \frac{2m(r)}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega^2 \ , \qquad (6)$$

The function $m(r)$ contains both the classical mass and a first-order quantum perturbation,

$$m(r) = M[1 + \epsilon\mu(r)] \ . \qquad (7)$$

The metric perturbation functions, $\rho(r)$ and $\mu(r)$, are determined by solving the semiclassical Einstein equations expanded to first order in ϵ,

$$\frac{d\mu}{dr} = -\frac{4\pi r^2}{M\epsilon}\langle T^t_{\ t}\rangle \ , \qquad (8)$$

$$\frac{d\rho}{dr} = \frac{4\pi r}{\epsilon}\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1}\left[\langle T^r_{\ r}\rangle - \langle T^t_{\ t}\rangle\right] \ . \qquad (9)$$

Assuming the perturbation expansion remains valid, the functions $\mu(r)$ and $\rho(r)$, obtained by integrating Eqs. (8-9), will contain constants of integration. It is convenient to define them as the values of the metric perturbations on the unperturbed horizon at r_+, so that $\mu(r_+) = C_1$ and $\rho(r_+) = C_2$. Since we are working in perturbation theory, the values of these quantities on the actual horizon are, to leading order, also C_1 and C_2 respectively. Then to first order in ϵ the value of $m(r)$ at the horizon is $m(r_h) = M(1 + \epsilon C_1)$. It is clear that C_1 represents a finite renormalization of the mass M of the black hole. As in previous work [3,4], we hereafter denote the renormalized perturbed mass at the horizon, $m(r_h) = M(1 + \epsilon C_1)$, by M_R.

The location of the horizon is given by the vanishing of $f(r_h)$ and $h^{-1}(r_h)$. This condition is equivalent to

$$r_h^2 - 2M_R r_h + Q^2 = 0 \ . \qquad (10)$$

Thus for a given value of r_h

$$M_R = \frac{1}{2}\left(r_h + \frac{Q^2}{r_h}\right) \ . \qquad (11)$$

There exists a sequence of black hole solutions which have $M_R = |Q| + O(\epsilon^2)$. For solutions in this sequence $r_h = |Q|(1 + \epsilon\, b)$ with b a constant of order unity or less.

The surface gravity for these solutions to first order in ϵ is

$$\kappa = \frac{\epsilon}{|Q|}\left[b - |Q|\mu'(|Q|)\right] \ . \qquad (12)$$

Thus as the value of r_h decreases for solutions in this sequence the surface gravity does also. The sequence terminates with a zero temperature black hole at

$$b = |Q|\,\epsilon\mu'(r_+) \ . \qquad (13)$$

It is the zero temperature solution found by Lowe [7]. From Eq. (8) one sees that $\mu'(|Q|)$ is proportional to the energy density of the quantized fields evaluated at $r = |Q|$ in the background spacetime. Since the energy density of these fields in an ERN background spacetime is negative at $r = |Q|$ in all physically realistic cases for free fields of spin 0, 1/2, and 1 [5], it is clear that if a zero temperature black hole solution is to exist then $b < 0$. In our previous paper [5] we considered solutions with $b \geq 0$.

There is a problem for the solutions with $b < 0$, corresponding to $r_h < |Q|$, including the zero temperature solution found by Lowe. In a perturbation expansion the stress-energy tensor of the quantized fields is computed in the background spacetime, and is only known for values $r \geq r_+$, where r_+ is the radius of the event horizon in the background (unperturbed) spacetime. But the event horizons of these proposed perturbative solutions lie inside the event horizons of the unperturbed RN or ERN black holes, $r_h < |Q| \leq r_+$. In particular for the ERN black hole there is absolutely no justification to extrapolating the stress-energy values inside r_+ because the event horizon is also a Cauchy horizon. Thus perturbation theory cannot be applied in these cases.

It is still conceivable that these solutions do represent a valid set of approximate solutions to the full nonlinear semiclassical equations. To establish this, however, it would be necessary to compute $\langle T_\mu^\nu \rangle$ in the semiclassical spacetime with $r_h < |Q|$ and check the self-consistency of the solution.

There is another potential problem that could occur for any proposed semiclassical solution with $\kappa = T = 0$. It has been shown by Trivedi [9] that in two spacetime dimensions the stress-energy tensor for the quantized scalar field diverges on the event horizons of most zero-temperature black holes including the (2-d) ERN black hole. In four dimensions two of us [8] have shown numerically that no such divergence exists for massless quantized scalar fields in the ERN background. However, it is quite possible that such a divergence could occur on the event horizon of the perturbed zero temperature black hole.

ACKNOWLEDGMENTS

P. R. A. would like to thank G. Cook, W. Kerr and O. Zaslavskii for helpful conversations. This work was supported in part by National Science Foundation Grant No. PHY-9734834 at Montana State University and Nos. PHY-9800971 and PHY-0070981 at Wake Forest University.

REFERENCES

* electronic mail address: anderson@wfu.edu
† electronic mail address: hiscock@physics.montana.edu
‡ electronic mail address: brett@peloton.runet.edu

[1] A. Strominger and C. Vafa, Phys. Lett. **B379**, 99 (1996).
[2] J. M. Bardeen, Phys. Rev. Lett. **46**, 382 (1981).
[3] J. W. York, Jr., Phys. Rev. D **31**, 775 (1985); D. Hochberg, T. W. Kephart, and J. W. York, Jr., Phys. Rev. D **48**, 479 (1993); P. R. Anderson, W. A. Hiscock, J. Whitesell, and J. W. York Jr., Phys. Rev. D **50**, 6427 (1994).
[4] B. E. Taylor, W. A. Hiscock, and P. R. Anderson, Phys. Rev. D. **61**, 84021 (2000).
[5] P. R. Anderson, W. A. Hiscock, and B. E. Taylor, Phys. Rev. Lett. **85**, 2438 (2000).
[6] Throughout we use units such that $\hbar = c = G = k_B = 1$. Our conventions are those of C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
[7] D. A. Lowe, Preprint, gr-qc/0011053.
[8] P. R. Anderson, W. A. Hiscock, and D. J. Loranz, Phys. Rev. Lett. **74**, 4365 (1995).
[9] S. P. Trivedi, Phys. Rev. D**47**, 4233 (1993).